

## **EVEN AFTER PERCUTANEOUS CORONARY INTERVENTION OF ANGIOGRAPHICALLY SIGNIFICANT LESIONS, IVUS-DEFINED HIGH-GRADE STENOSES ARE COMMON. A BASELINE IVUS ANALYSIS FROM THE PROSPECT TRIAL**

i2 Poster Contributions

Georgia World Congress Center, Hall B5

Monday, March 15, 2010, 9:30 a.m.-10:30 a.m.

---

Session Title: Intravascular Diagnostics and Complex Lesions

Abstract Category: Intravascular Diagnostics

Presentation Number: 2503-510

---

Authors: *Akiko Maehara, Gary S. Mintz, Ecaterina Cristea, Alexandra J. Lansky, Roxana Mehran, George Dangas, Barry Templin, Wai-Fung Cheong, John McPherson, Bernard de Bruyne, Naim Farhat, Steven Marso, Jean Fajadet, Michael Foster, Patrick W. Serruys, Gregg W. Stone, Cardiovascular Research Foundation and Columbia University Medical Center, New York City, NY*

PROSPECT (Providing Regional Observations to Study Predictors of Events in the Coronary Tree) was a multicenter, multimodality imaging study designed to prospectively identify vulnerable plaque after treatment of all culprit lesions in pts presenting with acute coronary syndromes. The protocol specified that all 3 coronary arteries - both culprit and non-culprit lesion containing vessels - be studied with angiography and intravascular ultrasound (IVUS) after successful PCI, including imaging of the proximal 6-8cm of each major epicardial artery with IVUS. Of 615 pts with analyzable IVUS, the average number of imaged vessels was  $2.84 \pm 0.50$  per pt, the average number of analyzable vessels was  $2.57 \pm 0.64$ , and the average length of analyzable arteries measured  $193 \pm 82$  mm. An IVUS lesion was defined as a plaque burden (plaque/vessel area)  $>40\%$  in 3 consecutive frames ( $\sim 1.5$  mm in length).

**Results:** 30.3% of pts presented with STEMI, 65.6% of pts presented with NSTEMI, and 4.4% with unstable angina and ECG changes. After treating all angiographically significant culprit lesions (72% of pts had culprit lesions in 1 artery, and 28% had culprit lesions in 2 arteries), IVUS identified 2698 residual lesions (mean  $4.5 \pm 2.1$  lesions per pt). Among these, there were 512 lesions (mean  $0.83 \pm 1.08$  per pt or at  $\geq 1$  such lesion in 42.2% of pts) that had an IVUS minimum lumen area (MLA)  $<4.0$  mm<sup>2</sup>, a commonly used and ischemia-validated criterion for a significant stenosis. Importantly, among IVUS lesions with an MLA  $<4.0$  mm<sup>2</sup>, an angiographic diameter stenosis  $>50\%$  was seen in only 2%. Finally, among 846 lesions detected by both angiography (visual diameter stenosis  $>30\%$ ) and IVUS, 251 angiographic lesions (29.7%) had an IVUS MLA  $<4.0$  mm<sup>2</sup> while 259 angiographic lesions (33.0%) had insufficient plaque to reach the criteria for an IVUS lesion ( $<40\%$  plaque burden or  $<3$  frames).

**Conclusions:** Even after treating all culprit lesions in pts with STEMI and NSTEMI, IVUS identified angiographically silent, but potentially clinically important stenoses in  $\approx 40\%$  of pts. Conversely, many suspected angiographic lesions did not have even modest plaque burden detected by IVUS.